

## Pediatric and Adolescent Health Partners Well Visit Guidelines 2018

At your child's well visits these activities are planned, depending upon his/her age:

1. Measurements: Height and Weight and Blood Pressure, Growth Chart Plotting with BMI
2. History of Well Being and Ongoing Problems with a Review of the Growth Chart
3. Functional testing: vision and hearing
5. Physical Examination
5. Laboratory testing: urine, blood or others as indicated
6. Child Care Advice: Nutrition, Safety, Cognitive- Emotional-Social Development
7. Management advice and prescriptions for ongoing problems
8. Immunizations
9. Completion of forms that you provide [Daycare, Preschool, School, Sports, Camps, etc.]

AGE	LAB/SCREENING	TEST	IMMUNIZATION
2 weeks			
6 weeks			(DTaP, Hib, IPV)= PENTACEL* Rotateq (oral), Prevnar -13
3 months			(DTaP, Hib, IPV)= PENTACEL Rotateq (oral), Prevnar -13
4 ½ months			Rotateq (oral), Prevnar-13, Hep B
6 months			(DTaP, Hib, IPV)= PENTACEL Hep B
9 months	Hemoglobin(Hgb)		Hep B
12 months	TB Risk		Varicella, Prevnar-13
15 months			MMR, Hep A**
18 months			(DTaP, Hib, IPV)= PENTACEL
2 years			Hep A* & catch-up any missed
3 years	Urine (if able)	Vision	Varicella, MMR
4 years	Urine	Vision	DTaP, IPV
5 years	Urine, TB Risk Hgb & Lead	Vision, Hearing	
6-10 years	Urine	Vision	
11 -14 years	Urine	Vision	Meningococcal (=Menactra), Tdap (=ADACEL) HPV*** (3 shot series=GARDASIL)
15-21 years	Urine	Vision	Catch-up if any shot is missed

\*Immunization schedule may vary depending upon your child, interfering illness, insurance coverage, vaccine availability, and changing CDC/ AAP recommendations.

\*\* Hepatitis A vaccine- now recommended for all children and adults in the USA.

\*\*\* HPV = Human Papillomavirus Vaccine: is available for both BOYS and GIRLS. Used for prevention of Cervical Cancer, Penile Cancer and Venereal Wart caused by HPV strains, it is FDA approved for ages 9-26, preferably before any sexual activity. We strongly recommend it.