

## Schedule of immunizations, labs, and screenings at well visits

AGE	LABS	SCREENING	IMMUNIZATIONS*
2 weeks		Edinburgh PPDS	
6 weeks		Edinburgh PPDS	Pentacel (DTaP,Hib,IPV), Pevnar 13, Rotateq (oral)
3 months		Edinburgh PPDS	Pentacel, Pevnar 13, Rotateq (oral)
4 ½ months			Hepatitis B, Rotateq (oral)
6 months			Pentacel, Pevnar 13
9 months			Hepatitis B
12 months	CBC (Hgb), lead screen, TB risks	fluoride varnish (if desired)	MMR, Pevnar 13
15 months			Pentacel, Varicella
18 months		M-CHAT, fluoride varnish (if desired)	Hepatitis B
2 years			Hepatitis A**
2 ½ years		M-CHAT, fluoride varnish (if desired)	Hepatitis A**
3 years	urinalysis	BP, vision	catch up as needed
4 years	urinalysis	BP, vision	DTaP, MMR
5 years	Hgb, lead screen, urinalysis, TB risks	BP, hearing, vision	IPV, Varicella
6-10 years	urinalysis	BP, vision	catch up as needed
11 years	urinalysis	BP, hearing, vision	Meningococcal (Menactra), Tdap (Adacel), (HPV information sheet)
12-14 years	urinalysis	BP, suicide screening, vision	HPV (Gardasil): 2 injections separated by 6-12 months if started prior to 15 <sup>th</sup> birthday
15-21 years	chlamydia screen, urinalysis	BP, hearing, suicide screening, vision	HPV(Gardasil): 3 injections over 6 months if started after the 15 <sup>th</sup> Birthday Meningococcal (Menactra) (booster after 16 <sup>th</sup> bday), Meningococcal B (Bexsero) (2 Total)

\*Influenza (flu) vaccination: recommended yearly starting at 6 months. First vaccination requires a booster 28 days after the first dose if receiving a flu vaccine for the first time and current age less than 9 years.

\*Immunization schedule may vary depending upon your child, interfering illness, insurance coverage, vaccine availability, and changing CDC/AAP recommendations.

\*\*Hepatitis A vaccine: Required in Western USA and sometimes needed for foreign travel. It is recommended for everyone older than 12 months.